



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# *Note on C. S. Peirce's Paper on "A Quincuncial Projection of the Sphere."*

BY JAMES PIERPONT, *New Haven, Conn.*

In the second volume of this journal\* Professor C. S. Peirce called attention to a very elegant representation of the sphere on the plane by means of the function  $\text{cn}\left(z, \kappa = \frac{1}{\sqrt{2}}\right)$ . Let  $\theta, l, p$  be the longitude, latitude, and north polar distance resp. of a point  $P$  on the sphere. If  $\zeta = \xi + i\eta$  be the stereographic projection of this point on the equatorial plane ("ζ-plane"), we have

$$\zeta = \rho e^{i\theta} = \tan \frac{p}{2} \cdot e^{i\theta}.$$

Let now  $\zeta = \text{cn}\left(z, \frac{1}{\sqrt{2}}\right)$ ; the ζ-plane and thus the sphere itself is conformally represented on the "z-plane." Being given ζ, it is not difficult to find formulæ for determining the coordinates of  $z$  and thus follow the movements of  $P$  in the  $z$ -plane.

The formulæ given by Prof. Peirce for this purpose are

$$x_\kappa = \frac{1}{2} F(\phi), \tag{1}$$

where  $x_\kappa$  is one of the coordinates of  $z = x + iy$ ,

$$\cos^2 \phi = \frac{\sqrt{1 - \cos^2 l \cos^2 \theta} - \sin l}{1 + \sqrt{1 - \cos^2 l \cos^2 \theta}}, \tag{2}$$

and as usual

$$F(\phi) = \int_0^\phi \frac{d\phi}{\sqrt{1 - \frac{1}{2} \sin^2 \phi}}.$$

There seems to be an error in this determination, however, as may be seen in

---

\*C. S. Peirce, "A Quincuncial Projection of the Sphere." *American Journal of Mathematics*, Vol. II (1879), p. 394.

taking special values of  $l$  and  $\theta$ . For example, for  $l = \theta = 0$  we have  $\zeta = 1$ , whence  $z \equiv 0 \pmod{4K, 2K(1+i)}$ , so that

$$x \equiv y \equiv 0. \quad (3)$$

The formulæ (1), (2), however, require that

$$x_\kappa \equiv \frac{K}{2} \pmod{K},$$

which contradicts (3).

Expressions for  $x, y$  may be determined as follows:\* Let  $u = x + iy$ ,  $v = x - iy$ ; then  $u + v = 2x$ ,  $u - v = 2iy$  and

$$\operatorname{cn} 2x = \frac{\operatorname{cn} u \operatorname{cn} v - \operatorname{sn} u \operatorname{sn} v \operatorname{dn} u \operatorname{dn} v}{1 - \frac{1}{2} \operatorname{sn}^2 u \operatorname{sn}^2 v}.$$

Now  $\operatorname{cn} u = \rho e^{i\theta}$ , whence  $\operatorname{cn} v = \rho e^{-i\theta}$ . Similarly let

$$\begin{aligned} \operatorname{sn} u &= \rho_1 e^{i\theta_1}, & \operatorname{dn} u &= \rho_2 e^{i\theta_2}, \\ \operatorname{sn} v &= \rho_1 e^{-i\theta_1}, & \operatorname{dn} v &= \rho_2 e^{-i\theta_2}, \end{aligned}$$

so that

whence

$$\operatorname{cn} 2x = \frac{\rho^2 - \rho_1^2 \rho_2^2}{1 - \frac{1}{2} \rho_1^4} \quad (4)$$

and

$$\operatorname{cn} 2y = \frac{1 - \frac{1}{2} \rho_1^4}{\rho^2 + \rho_1^2 \rho_2^2}. \quad (5)$$

But

$$\rho_1^4 = \frac{4(1 - \cos^2 \theta \cos^2 l)}{(1 + \sin l)^2}, \quad \rho_2^4 = \frac{1 - \sin^2 \theta \cos^2 l}{(1 + \sin l)^2}.$$

Thus introducing the angles  $\alpha, \beta$ , the equations (4), (5) give

$$\left. \begin{aligned} \operatorname{cn} 2x &= \frac{\cos^2 l - 2\sqrt{\sin^2 l + \frac{1}{4} \cos^4 l \sin^2 2\theta}}{2 \sin l + \cos^2 l \cos 2\theta} = \cos \alpha, \\ \operatorname{cn} 2y &= \frac{2 \sin l + \cos^2 l \cos 2\theta}{\cos^2 l + 2\sqrt{\sin^2 l + \frac{1}{4} \cos^4 l \sin^2 2\theta}} = \cos \beta, \end{aligned} \right\} \quad (6)$$

whence

$$x = \frac{1}{2} F(\alpha), \quad y = \frac{1}{2} F(\beta). \quad (7)$$

The corresponding formulæ expressing  $\xi, \eta$  in terms of  $x, y$  are

$$\xi = \frac{\operatorname{cn} x \operatorname{cn} y}{1 - \operatorname{sn}^2 y \operatorname{dn}^2 x}, \quad \eta = -\frac{\operatorname{sn} x \operatorname{sn} y \operatorname{dn} x \operatorname{dn} y}{1 - \operatorname{sn}^2 y \operatorname{dn}^2 x}. \quad (8)$$

Before computing the coordinates of  $z$  for a point  $P$  on the sphere, it is well to see what general correspondence exists between the  $z$ -plane and the sphere.

---

\* Cf. Richelot, "Darstellung einer beliebigen Grösse durch  $\sin am(u+w, k)$ ." *Crelle*, Vol. 45. Durège, "Theorie der elliptischen Functionen." 4th ed. Leipzig, 1887, p. 289.

Figure 1 represents the stereographic projection of the sphere on the  $\zeta$ -plane. The inner circle,  $\alpha$ , has a radius  $= 1$ ; the outer circle,  $\beta$ , has an infinite radius; to them correspond on the sphere resp. the equator and an infinitely small circle about the south pole. Points on the northern hemisphere are projected within  $\alpha$ , points on the southern hemisphere without  $\alpha$ .

The point  $\alpha_0$  represents the  $N$ -pole. The eight lines passing through  $\alpha_0$  and  $\beta_1, \beta_2, \dots$  represent the eight meridian circles whose longitudes are resp.  $\theta = 0^\circ, 45^\circ, 90^\circ, \dots$ . For shortness I shall designate any line by its terminal points; thus  $(\alpha_2 \alpha_4)$  denotes for example the arc of  $\alpha$  terminated by  $\alpha_2, \alpha_4$ . Similarly  $\{\dots\}$  shall represent a surface bounded by lines of the figure passing through the points within  $\{\}$ .

Let us now turn to the correspondence between the  $\zeta$ - and  $z$ -plane. The parallelogram  $\Pi = \{ABCD\}$  in Fig. 2 being an elementary parallelogram of periods of the function  $\text{cn}\left(z, \frac{1}{\sqrt{2}}\right)$ , a point of the  $\zeta$ -plane is represented by two points  $z_1, z_2$  in  $\Pi$ . Since  $z_1 + z_2 \equiv 0 \pmod{4K, 2K(1+i)}$ , the points  $z_1, z_2$  are symmetrical with respect to  $a_3$  as a center. This shows that  $\{DBC\}$  or  $\{DAB\}$  represents once and only once every point in the  $\zeta$ -plane, and conversely. Instead, however, of employing  $\Pi$  to represent the  $\zeta$ -plane, we may use the square  $\Sigma = \{A'B'C'D'\}$ . It consists of four lesser squares  $\Sigma_1 \Sigma_2 \Sigma_3 \Sigma_4$ , the first two having  $\alpha_0, i_2$  resp. for centers. As the point  $a_3$  is a center of symmetry (in the afore sense) the  $\zeta$ -plane is represented once only by  $\Sigma' = \Sigma_1 + \Sigma_2$ . Thus corresponding to a point on the sphere there exists one point only in  $\Sigma'$ , and conversely. Further to the  $N$ -hemisphere corresponds  $\Sigma_1$  and to the  $S$ -hemisphere corresponds  $\Sigma_2$ ; the equator is thus represented by the perimeters  $a, a'$  of  $\Sigma_1, \Sigma_2$ .

The point  $\alpha_0$  represents the  $N$ -pole,  $i_2$  the  $S$ -pole. In general, corresponding points in the two planes are marked by the same letters and suffixes; points in the  $\zeta$ -plane bearing Greek letters, and those of the  $z$ -plane, Latin. Finally, when  $\zeta$  moves on any line marked in Fig. 1,  $z$  moves in Fig. 2 on corresponding lines. Between the points of a surface  $\sigma$  enclosed by such a path of  $\zeta$  and the points of the corresponding surface  $s$  in the  $z$ -plane, there exists 1—1 correspondence. In particular, the squares  $\Sigma_1, \Sigma_2$  are divided into 16 triangles  $t$ , as e. g.  $\{\alpha_0 \alpha_1 \alpha_8\}$ , to each of which corresponds an  $\frac{1}{8}$  of a hemisphere. The proof of these statements I will illustrate for one or two cases.

For example, to see that when  $z$  describes  $(\alpha_0 \alpha_7)$ ,  $\zeta$  describes continuously




FIG. 1.— $\zeta$ -plane.




FIG. 2.— $z$ -plane.

and without returning on itself ( $\alpha_0 \alpha_7$ ), we set  $z = K + iy$ ,  $0 \leq y \leq K$ . Then

$$\zeta = \text{cn}(K + iy) = -\kappa \frac{\text{sn } iy}{\text{dn } iy} = -\iota \kappa \frac{\text{sn } y}{\text{dn } y}, \quad \kappa = \frac{1}{\sqrt{2}}.$$

For  $y = 0$ ,  $\zeta = 0$ ; for  $y = K$ ,  $\zeta = -i$ . As  $\zeta$  is a continuous function of  $y$ ,  $\zeta$  moves continuously along ( $\alpha_0 \alpha_7$ ); further, it cannot return on itself, for then a point  $\zeta$  on ( $\alpha_0 \alpha_7$ ) would be twice represented on ( $\alpha_0 \alpha_7$ ).

To establish the correspondence between  $\alpha$  and  $a$ ,  $a'$ , let us show for example that ( $a_1 \alpha_7$ ) corresponds to ( $\alpha_1 \alpha_7$ ). Here  $z = x(1 + i)$  and thus

$$\zeta = \text{cn}(x + ix) = \frac{\text{cn}^2 x - i \text{sn}^2 x \text{dn}^2 x}{1 - \text{sn}^2 x \text{dn}^2 x}, \quad \therefore |\zeta| = 1,$$

so that  $\zeta$  describes continuously and without turning the quadrant ( $\alpha_1 \alpha_7$ ) while  $z$  moves on ( $a_1 \alpha_7$ ).

In a similar manner we can establish the correspondence between the medial lines ( $\alpha_2 \alpha_6$ ), ( $\alpha_4 \alpha_8$ ) of the square  $\Sigma_1$  and the diameters ( $\alpha_2 \alpha_6$ ), ( $\alpha_4 \alpha_8$ ) of the circle  $\alpha$ . This can also be readily shown by employing (8). Their quotient, namely, gives

$$\tan \theta = - \frac{\text{sn } x \text{sn } y \text{dn } x \text{dn } y}{\text{cn } x \text{cn } y}.$$

As here  $x = y + K$ ,  $\tan \theta = 1$ , so that ( $a_2 \alpha_6$ ) corresponds to ( $\alpha_2 \alpha_6$ ). The correspondence of the circles  $b$ ,  $\beta$  is thus shown. For points in the vicinity of  $i_2$ ,  $z = \iota K + z'$  and

$$\begin{aligned} \zeta &= \text{cn}(\iota K + z') = -\frac{\iota \text{dn } z'}{\kappa \text{sn } z'} \\ &= \frac{1}{z'} (a + bz'^2 + \dots), \end{aligned}$$

which shows that while  $z$  describes a small circle in positive sense about  $i_2$ ,  $\zeta$  describes an infinitely large circle in negative sense. An inspection of the relative position of the triangles  $t$ ,  $\tau$  leads one to suspect that the diagonals and medial lines of  $\Sigma_1$ ,  $\Sigma_2$  are lines of symmetry; that is, if  $z_1$ ,  $z_2$  be two points in  $z$ -plane situated symmetrically in respect to such a line, then  $\zeta_1$ ,  $\zeta_2$  are symmetrical with respect to the corresponding line in the  $\zeta$ -plane. That this is so can be illustrated on the medial ( $\alpha_2 \alpha_6$ ). For if in (8) we replace  $x$  by  $K + y$  and  $y$  by  $x - K$ , the expressions for  $\xi$ ,  $\eta$  interchange. For example, the expression

for  $\xi$  becomes

$$\frac{\operatorname{cn}(K+y)\operatorname{cn}(x-K)}{1-\operatorname{dn}^2(K+y)\operatorname{sn}^2(x-K)} = x^2 \frac{\operatorname{sn} x \operatorname{sn} y \operatorname{dn} x \operatorname{dn} y}{\operatorname{dn}^2 x \operatorname{dn}^2 y - x^2 \operatorname{cn} x} = \eta.$$

In the same way we show that the diagonal  $(a_3 a_7)$  is an axis of symmetry. For if  $z_1 = K + a + ib$  and  $z_2 = K - a + ib$  be two such points, we have

$$\begin{aligned} \zeta_1 &= \operatorname{cn}(K + a + ib) = -x \frac{\operatorname{sn}(a + ib)}{\operatorname{dn}(a + ib)} \\ &= -x \frac{\operatorname{sn} a \operatorname{dn} b + i \operatorname{sn} b \operatorname{cn} a \operatorname{cn} b \operatorname{dn} a}{\operatorname{cn} b \operatorname{dn} a \operatorname{dn} b - x^2 \operatorname{sn} a \operatorname{sn} b \operatorname{cn} a} = \frac{A + iB}{C - iD} \\ &= \frac{AC - BD + i(BC + AD)}{C^2 + D^2} = \xi + i\eta. \end{aligned}$$

Replacing  $a$  by  $-a$ ,  $A$  and  $D$  become  $-A$ ,  $-D$  and

$$\zeta_2 = -\xi + i\eta.$$

We may form a good idea of the representation of the parallels and meridians in the  $z$ -plane by considering the expression for the magnification

$$m = \left| \frac{dz}{dp} \right| = \left| \frac{dz}{d\zeta} \cdot \frac{d\zeta}{dp} \right|.$$

As  $\frac{d\zeta}{dp} = \frac{1}{1 + \sin l}$  and  $\frac{dz}{d\zeta} = \frac{1}{\rho_1 \rho_2}$ ,

$$m = \frac{1}{\sqrt{2}} \frac{1}{\sqrt{\sin^2 l + \frac{1}{4} \cos^2 l \sin^2 2\theta}}.$$

This shows that the magnification  $m$  is greatest at the equator and least at the poles; also that along a parallel  $m$  has a minimum for  $\theta = 45^\circ, 135^\circ \dots$  and a maximum for  $\theta = 0^\circ, 90^\circ \dots$ . We have already seen that the representation is conform in the vicinity of the  $S$ -pole. The same being true for the  $N$ -pole, the parallels are approximately represented in the  $z$ -plane for some distance from the poles by circles. As however they approach the equator, the above considerations show that they take on square-like forms with rounded corners. As the representation is in general conform, the  $z$ -plane meridians everywhere cut the just described parallels at right angles, so that as they depart from the lines  $(a_2 a_6)$ , etc., they bend inward toward the same.

The correspondence of the planes  $\zeta$  and  $z$  being now pretty accurately established, we may employ the formulæ (6), (7) with advantage. The foregoing considerations show that we need to compute  $\alpha, \beta$  only for values of  $\theta$  lying

between  $0^\circ$  and  $45^\circ$ . We may, if we like, use only one of the angles  $\alpha, \beta$ , in which case we must take  $\theta$  between  $0^\circ$  and  $90^\circ$ . As an illustration of their use I append the following table for  $l = 5^\circ$ :

$\theta$	$\alpha$	$\beta$	$x$	$y$	$\frac{x}{K}$	$\frac{y}{K}$	$1 - \frac{x}{K}$
0	$45^\circ 29$	0	.4181	0	.2255	0	.7745
5	49 32	$21^\circ 28$	.4593	.1895	.2477	.1022	.7523
15	63 10	47 5	.6065	.4341	.3271	.2124	.6729
45	85 0	85 0	.9887	.8654	.5333	.4668	.4667

It will be noticed that although the formula given by Prof. Peirce for computing the coordinates of  $z$  is incorrect, the last two columns of the above table agree with the results given by him.

The representation afforded by  $\zeta = \text{cn} \left( z, \frac{1}{\sqrt{2}} \right)$  is everywhere conform except for certain points for which  $\frac{d\zeta}{dz}$  becomes 0 or  $\infty$ , that for the corners of  $\Sigma_1, \Sigma_2$  and  $i_2$ . We have already seen that the representation is conform at  $i_2$ , a fact illustrated by the diagonals and medial lines of  $\Sigma_2$ .

For the other points, however, the representation is not conform. To take an example  $a_1$ . For its vicinity,  $\zeta = \text{cn } z = (1 + az^2 + bz^4 + \dots)$ , and thus

$$\zeta - 1 = z^2(a + bz^2 + \dots),$$

which shows that two lines in the  $z$ -plane meeting under the angle  $\phi$  meet under the angle  $2\phi$  in the  $\zeta$ -plane. Thus  $(a_1 a_2), (a_1 a_6)$  make an angle of  $90^\circ$  in the  $z$ -plane, while in the  $\zeta$ -plane they meet under an angle of  $180^\circ$ . Similarly  $(a_1 a_6), (a_1 a_2)$  meet under the angle  $45^\circ$  in  $z$ -plane, and under the angle  $90^\circ$  in the  $\zeta$ -plane. At all the corners of  $\Sigma_1, \Sigma_2$  the distortion of angles is double.

The function  $\zeta = \text{cn} \left( z, \frac{1}{\sqrt{2}} \right)$  possesses then the very remarkable property



of representing in 1—1 correspondence the interior of the square  $\Sigma_1$  by the interior of a circle of unit radius about the origin of the  $\zeta$ -plane. Only at the corners does this representation cease to be conformal.

That such a function existed was discovered by Schwarz\* while searching to determine a function, under certain simple conditions, to illustrate Riemann's theorem† that it is possible in one way only to represent conformally a simply connected surface  $T$  on a circle so that to the center corresponds any point in the interior, and to a point on the circumference any point on the edge of  $T$ . For the case of a square whose corners were  $\pm K, \pm \iota K$ , Schwarz arrived at the function  $\zeta = \text{sn}(u, \iota)$ , which may also be written  $\zeta = \text{cn}\left(z, \frac{1}{\sqrt{2}}\right)$ , where  $z = K - \sqrt{2}u$ . This relation enables us to deduce all properties of  $\text{cn}\left(z, \frac{1}{\sqrt{2}}\right)$  immediately from those of  $\text{sn}(u, \iota)$ , or conversely.

\*Schwarz, Crelle, vol. 70 (1869), p. 105-120; also *Gesam. Math. Abh.*, vol. II, p. 65: Ueber einige Abbildungsaufgaben.

† Riemann, *Gesam. Werke*, p. 40. Grundlagen für eine allgemeine Theorie der Functionen einer complexen Veränderlichen.